
Use of the Free Electron Laser for the Noninvasive Determination of Retinal Oxyhemoglobin Saturation by Near Infrared Reflectance Spectrophotometry














2000 National Estimates of the Burden of Diabetes

Diabetes

Prevalence:

Diagnosed Undiagnosed

Total

11.1 million

5.9 million


17.0 million (6.2% of US population)

Diabetic Retinopathy Prevalence: 65,000 diabetics per year develop proliferative DR


Leading cause of blindness among 20 to 74 yr old


Ref: National Diabetes Information Clearinghouse Research to Prevent Blindness
























Backscatter


Favorable Attributes of the FEL


- Short pulsation of the emitted light
 - » Minimizes total retinal irradiance
 - » Compensates for eye movements














Optical Absorption of Hemoglobin



Optical Absorption of Hemoglobin




Optical Absorption of Hemoglobin





Favorable Attributes of the FEL

- Broad tunability of the emitted light.
 - » To rapidly scan a range of 70 to 100 nm
 - » Alternate between visible and near-infrared
- Strong energy output in the near-infrared region of the spectrum (700 to 900 nm).
 - Where hemoglobin chromophore signal is attenuated












Derivation of Algorithm


Consulted References

- •R. Pittman and B. Duling (1975) described the relationship between spectral absorption and oxygen saturation of hemoglobin in microvasculature within visible range.
 - •A.P. Sheperd (1975) developed an empirically derived analytical expression relating OD and Hb concentration.
- •Delori (1988) Determined contribution of fundus scattering to OD of hemoglobin at three wavelengths (555nm).
- •Ferrari et al (1989) utilized DNIRS to noninvasively determine cerebral venous Hb saturation (sagittal sinus).











Ref: Padnick-Silver, Linsenmeier RA. IOVS 44(2):745-750, 2003. Linsenmeier RA. J. Gen. Physiol. 88:521-542, 1986

In Situ Schematics of Optical System







Summary

- Presents the opportunity to study retinal metabolism in a yet unprecedented fashion.
- •Technique may reveal subtle alterations in retinal metabolism that are not discernable with present routine function tests.
- •Could promote a more rational physiologic basis for clinical intervention and delve into the pathogenic mechanisms and possible early diagnosis of diabetic retinopathy, and other retinal ischemic/vascular diseases.
- Collectively these retinopathies comprise the leading causes of blindness in the industrialized world.

